

AD-A175 397

CLONING OF A CDNA ENCODING THE CALMODULIN BINDING
DOMAIN OF THE PLASMA ME. (U) KENTUCKY UNIV LEXINGTON
DEPT OF BIOCHEMISTRY P BRANDT 17 NOV 86

1/1


UNCLASSIFIED

ARO-20358.1-L5-F DARG29-83-G-0006

F/G 6/3

NL






MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

②

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARO 20358-1-45-F	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) CLONING OF A cDNA ENCODING THE CALMODULIN BINDING DOMAIN OF THE PLASMA MEMBRANE Ca^{2+} -ATPase		5. TYPE OF REPORT & PERIOD COVERED Final (8/15/83-8/14/86)
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Paul Brandt		8. CONTRACT OR GRANT NUMBER(s) DAAG29-83-G-0006
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Biochemistry University of Kentucky Lexington, KY 40536-0084		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE 11/17/86
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 5
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) calcium, calcium-ATPase, calmodulin, cDNA, lambda gt11, myosin light chain kinase.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A bacteriophage (Xgt11) cDNA expression library, derived from bovine brain mRNA, was screened with polyclonal antibodies raised against the plasma membrane calcium pumping ATPase (Ca^{2+} -ATPase). A positive phage was isolated and purified to homogeneity in three subsequent rescreenings. The expressed protein product was shown to be reactive with antibodies against the calmodulin binding domain of the Ca^{2+} -ATPase. The cDNA sequence showed homology with the SR Ca^{2+} -ATPase 3'-end noncoding region and the		



DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

86 12 30 054

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A175 397

CONFIDENTIAL
UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

derived protein sequence bore homology with the protein sequences of myosin light chain kinase calmodulin binding domain from smooth and skeletal muscle.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CLONING OF A cDNA ENCODING THE CALMODULIN
BINDING DOMAIN OF THE PLASMA MEMBRANE Ca^{2+} -ATPase

Final Report

Paul Brandt

Nov. 17, 1986

U.S. Army Laboratory Command-Army Research Office
Research Triangle Park, North Carolina

DAAG29-83-G-0006

Department of Biochemistry
University of Kentucky College of Medicine
Lexington, Kentucky

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



FINAL REPORT OF RESEARCH FINDINGS

Introduction

The divalent cation, Ca^{2+} , acts as a major second messenger for numerous processes in biological systems. Studies over the past 15 years have provided a substantial understanding of the molecular bases through which calcium signals act via intracellular receptors such as calmodulin (CaM) as has been summarized in numerous recent reviews (see for example, Klee and Vanaman, 1982; Veigl et al., 1984). However, the ability of the cell to use free Ca^{2+} concentration as a regulatory signal results from the fact that resting state concentrations of this cation are normally maintained at below micromolar concentrations by the action of a set of sequestration systems found in all cell types. On stimulation, cytosolic Ca^{2+} levels rise transiently via influx and release from internal stores to concentrations of 10^{-5} to 10^{-4} M to give increases in the activities of those enzymes involved in mounting the required tissue specific response. Termination of this response occurs when Ca^{2+} -levels are returned to the resting state levels. While bulk transport of Ca^{2+} is accomplished by a Ca^{2+} - Na^{+} exchange protein, this system only brings the intracellular Ca^{2+} levels to about 10^{-5} M. The plasma membrane Ca^{2+} pumping ATPase (Ca^{2+} -ATPase) is then utilized to reach the resting level.

The major form of the Ca^{2+} -ATPase found in all cell types is that found in the plasma membrane. To date, only the human erythrocyte membrane enzyme has been studied in detail. As set forth in the following sections, this enzyme is activated by CaM linking its activity to cellular responses. Substantial work is currently in progress in this and other laboratories to elucidate the structural features of this enzyme which specify membrane association, ATP hydrolysis, cation translocation and CaM regulation. Traditional techniques of protein chemistry applied to the purified ATPase have been the only approach to such studies to date. The large size of this enzyme (138 kDa) makes such studies difficult and precludes determination of the entire structure of the enzyme. We have recently developed a series of immunoreagents which are suitable for use in attempts to clone the ATPase. This would greatly facilitate structural studies as well as provide the necessary probes for examining its structural gene, its expression in different cell types, its distribution throughout animal species and possible alteration of its structure or expression in various disease states.

Final Report

We have successfully cloned a portion of the Ca^{2+} -ATPase of plasma membranes. This was done by screening a cDNA library inserted into a bacteriophage vector (λ gt11) that produces a fusion protein of β -galactosidase and the inserted cDNA. The screening was done with a polyclonal antibody that had been raised in rabbits against human erythrocyte Ca^{2+} -ATPase and was affinity purified by passing whole immune serum over a column containing covalently bound Ca^{2+} -ATPase. The Ca^{2+} -ATPase specific antibody was then eluted with a low pH buffer. The λ gt11 cDNA library was made from bovine brain messenger RNA by standard cDNA synthesis techniques.

As noted, the cloning vector used in these experiments yields an expressed protein that is coded for by the inserted cDNA. This allowed us to test the validity of a putative positive bacteriophage in a number of ways.

First, a cellular extract made from bacteria infected with the bacteriophage believed to contain a cDNA insert coding for a portion of the Ca^{2+} -ATPase was resolved by denaturing polyacrylamide gel electrophoresis and transferred to nitrocellulose paper. The expressed protein was detected by Ca^{2+} -ATPase specific immunostaining and was shown to be about 130kDa in size. This is to be expected since the fusion protein will be the sum of the molecular weights of the β -galactosidase and the Ca^{2+} -ATPase fragment coded for by the cDNA. This piece of data shows that the isolated cDNA codes for a 16 kDa piece of the Ca^{2+} -ATPase, or about 20% of the intact protein. Additional evidence shows that antibodies purified against the fusion protein react with purified Ca^{2+} -ATPase. In particular, these antibodies react only with fragments generated by a limited tryptic digest of the Ca^{2+} -ATPase known to contain the calmodulin binding domain (Zurini, et al., 1984). When these antibodies are added to a Ca^{2+} -ATPase assay they give partial stimulation of the Ca^{2+} -ATPase but not complete stimulation to the level obtained when calmodulin is used. These antibodies also block further activation by calmodulin to the Ca^{2+} -ATPase's normal level of stimulation. These data indicate that we have successfully cloned a cDNA coding for the calmodulin binding domain of the plasma membrane Ca^{2+} -ATPase.

Using the Sanger dideoxy sequencing method we have sequenced the entire cDNA. We have found that the cDNA is 1456 bp long and codes for 93 amino acids. There are 1177 bp of 3'-end noncoding region. Protein sequence homology comparisons with the calmodulin binding domains of skeletal muscle myosin light chain kinase (MLCK) (Blumenthal, et al., 1985) and smooth muscle MLCK (A. Means, personal communication) show regions of strong homology. These regions are conserved in both MLCK sequences, again suggesting we have isolated a cDNA encoding the calmodulin binding domain of the Ca^{2+} -ATPase. Comparison at the DNA level with the 3'-end of the sarcomplasmic reticulum (SR) Ca^{2+} -ATPase shows homology. Interestingly, this homology follows into the coding sequence of the plasma membrane (PM) Ca^{2+} -ATPase, but falls off when the sequence coding for the calmodulin binding domain is encountered. The SR Ca^{2+} -ATPase is not stimulated by calmodulin.

We have, therefore, isolated a cDNA fragment coding for a portion of the Ca^{2+} -ATPase from bovine brain. The validity of this fragment coding for Ca^{2+} -ATPase has been confirmed by several tests based on reactivity of a protein coded for by the cDNA with antibodies against purified Ca^{2+} -ATPase. Also, the protein sequence deduced from cDNA sequence is homologous to other calmodulin binding domains isolated from other calmodulin stimulated enzymes. The cDNA sequence is also homologous to the SR Ca^{2+} -ATPase cDNA sequence.

References

- Blumenthal, D.K., Takio, K., Edelman, A.M., Charbonneau, H., Titani, K., Walsh, K.A. and Krebs, E.G. 1985. Proc. Natl. Acad. Sci. 82:3187.
- Klee, C.B. and Vanaman, T.C. 1982. Adv. Prot. Chem. 35:213.
- MacLennan, D.H., Brandl, C.J., Korczak, B. and Green, N.M. 1985. Nature 316:696.
- Veigl, M.L., Vanaman, T.C. and Sedwick, W.D. 1984. BBA Reviews in Cancer 738:21.
- Zurini, M., et al. 1984. J. Biol. Chem. 259:618.

List of Publications

Isolation and Characterization of a Recombinant Phage Containing a Region of the Plasma Membrane Calcium Pumping ATPase cDNA. P. Brandt, M. Zurini, R.E. Rhoads, T.C. Vanaman, Fed. Proc. 1986. 45:1695.

Isolation of a cDNA Clone Coding for the Calmodulin Binding Domain of the Plasma Membrane Calcium Pumping ATPase. P.Brandt, M. Zurini. R. Neve, R.E. Rhoads, T.C. Vanaman, Proc. Natl. Acad. Sci. (in preparation).

Scientific Personnel Supported by DAAG29-83-G-0006

Paul Brandt - Graduate Student

END

2-87.

DTIC